
References

- Addario-Berry, L., Chor, B., Hallett, M., Lagergren, J., Panconesi, A. and Wareham, T. (2004). Ancestral maximum likelihood of phylogenetic trees is hard. *J. Bioinf. Comp. Biol.* **2**: 257–271.
- Aguinaldo, A.M., Turbeville, J.M., Linford, L.S., Rivera, M.C., Garey, J.R., Raff, R.A. and Lake, J.A. (1997). Evidence for a clade of nematodes, arthropods and other moulting animals. *Nature* **387**: 489–493.
- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In *Second International Symposium on Information Theory*, Tsahkadsor, Armenia, USSR, September 2–8, 1971 (eds B.N. Petrov and F. Csaaki), pp. 267–281. Budapest, Akademiai Kiado.
- Albert, V.A. and Mishler, B.D. (1992). On the rationale and utility of weighting nucleotide sequence data. *Cladistics* **8**: 73–83.
- Albert, V.A., Mishler, B.D. and Chase, M.W. (1992). Character-state weighting for restriction site data in phylogenetic reconstruction, with an example from chloroplast DNA. In *Molecular Systematics of Plants*. (eds P.S. Soltis, D.E. Soltis, and J.J. Doyle), pp. 369–403. New York, Chapman and Hall.
- Albert, V.A., Chase, M.W. and Mishler, B.D. (1993). Character-state weighting for cladistic analysis of protein-coding DNA sequences. *Ann. Missouri Bot. Gard.* **80**: 752–766.
- Albert, V.A., Backlund, A., Bremer, K., Chase, M.W., Manhart, J.R., Mishler, B.D. and K.C. Nixon. (1994). Functional constraints and *rbcL* evidence for land plant phylogeny. *Ann. Missouri Bot. Gard.* **81**: 534–567.
- Alfaro, M., Zoller, S. and Lutzoni, F. (2003). Bayes or bootstrap? A simulation study comparing the performance of Bayesian Markov chain Monte Carlo sampling and bootstrapping in assessing phylogenetic confidence. *Mol. Biol. Evol.* **20**: 255–266.
- Altschul, S.F. (1989). Gap costs for multiple alignment. *J. Theor. Biol.* **138**: 297–309.
- Aravind, L., Watanabe, H., Lipman, D.J. and Koonin, E.V. (2000). Lineage-specific loss and divergence of functionally linked genes in eukaryotes. *Proc. Natl. Acad. Sci. USA* **97**: 11319–11324.
- Ariew, A. (1998). Are probabilities necessary for evolutionary explanations? *Biol. Philos.* **13**: 245–253.
- Arvestad, L., Berglund, A.C., Lagergren, J. and Sennblad, B. (2003). Bayesian gene/species tree reconciliation and orthology analysis using MCMC. *Bioinformatics* **19** Suppl. 1: i7–i15.
- Avise, J.C. (1989). Gene trees and organismal histories: a phylogenetic approach to population biology. *Evolution* **43**:1192–1208.
- Avise, J.C. (2000). *Phylogeography: The History and Formation of Species*. Cambridge, MA, Harvard University Press.
- Bach, E. (1981). On time, tense and aspect: An essay in English metaphysics. In *Radical Pragmatics* (ed. P. Cole), pp. 63–81. New York, Academic Press.
- Baker, A. (2003). Quantitative parsimony and explanatory power. *Br. J. Phil. Sci.* **54**: 245–259.
- Bandelt, H.J., Forster, P., Sykes, B.C. and Richards, M.B. (1995). Mitochondrial portraits of human populations using median networks. *Genetics* **141**: 743–753.
- Barnes, E.C. (2000). Ockham's Razor and the anti-superfluity principle. *Erkenntnis* **53**: 353–374.
- Barrett, M., Donoghue M.J. and Sober, E. (1991). Against consensus. *Syst. Zool.* **40**: 486–493.
- Barry, D. and Hartigan, J.A. (1987). Statistical analysis of hominoid molecular evolution. *Stat. Sci.* **2**: 191–210.
- Benner, S.A., Trabesinger, N. and Scheiber, D. (1998). Post-genomic science: Converting primary sequence into physiological function. *Adv. Enzyme Regul.* **38**: 155–190.
- Benner, S.A., Chamberlin, S.G., Liberles, D.A., Govindarajan, S. and Knecht, L. (2000). Functional inferences from reconstructed evolutionary biology involving rectified databases — an evolutionarily grounded approach to functional genomics. *Res. Microbiol.* **151**: 97–106.
- Benson, D.A., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J. and Wheeler, D.L. (2004). GenBank: update. *Nucleic Acids Res.* **32**: D23–D26.
- Bi, S., Garilova, O., Gong, D.W., Mason, M.M. and Reitman, M. (1997). Identification of a placental enhancer for the human leptin gene. *J. Biol. Chem.* **272**: 30583–30588.

- Blackburn, D.G. (1984). From whale toes to snake eyes: Comments on the reversibility of evolution. *Syst. Zool.* **33**: 241–245.
- Blair, J.E., Ikeo, K., Gojobori, T. and Hedges, S.B. (2002). The evolutionary position of nematodes. *BMC Evol. Biol.* **2**: 7.
- Bock, W.J. (1973). Philosophical foundations of classical evolutionary classification. *Syst. Zool.* **22**: 375–392.
- Boudet, N., Aubourg, S., Toffano-Nioche, C., Kreis, M. and Lecharny, A. (2001). Evolution of intron/exon structure of DEAD helicase family genes in *Arabidopsis*, *Caenorhabditis*, and *Drosophila*. *Genome Res.* **11**: 2101–2114.
- Boyd, R. (1991). Confirmation, semantics, and the interpretation of scientific theories. In *The Philosophy of Science* (eds R. Boyd, P. Gasper and J.D. Trout), pp. 3–35. Cambridge, MA, MIT Press.
- Brady, R.H. (1985). On the independence of systematics. *Cladistics* **1**: 113–126.
- Brady, R.H. (1994). Explanation, description, and the meaning of transformation in taxonomic evidence. In *Models in Phylogeny Reconstruction* (eds R.W. Scotland, D.J. Siebert and D.M. Williams), pp. 11–29 special vol. 52, Syst. Assoc. Oxford, Clarendon Press.
- Bremer, K. (1988). The limits of amino acid sequence data in angiosperm phylogenetic reconstruction. *Evolution* **42**: 795–803.
- Bremer, K. (1994). Branch support and tree stability. *Cladistics* **10**: 295–304.
- Brooks, D.R. (1981). Hennig's parasitological method: A proposed solution. *Syst. Zool.* **30**: 229–249.
- Brooks, D.R. (1988). Scaling effects in historical biogeography: A new view of space, time, and form. *Syst. Zool.* **37**: 237–244.
- Brower, A.V.Z. (2000). Homology and the inference of systematic relationships: Some historical and philosophical perspectives. In *Homology and Systematics: Coding Characters for Phylogenetic Analysis* (eds R. Scotland and R.T. Pennington), pp. 10–21. New York, Taylor and Francis.
- Brudno, M., Chapman, M., Götting, B., Batzoglou, S. and Morgenstern, B. (2003). Fast and sensitive multiple alignment of large genomic sequences. *BMC Bioinformatics* **4**: 66.
- Brudno, M., Poliakov, A., Salamov, A., Cooper, G.M., Sidow, A., Rubin, E.M., Solovyev, V., Batzoglou, S. and Dubchak, I. (2004). Automated whole-genome multiple alignment of rat, mouse, and human. *Genome Res.* **14**: 685–692.
- Bryant, D. (2000). A lower bound for the breakpoint phylogeny problem. In *Proceedings of the 11th Annual Symposium on Combinatorial Pattern Matching* (eds R. Giancarlo and D. Sankoff), pp. 235–247. London, Springer Verlag.
- Bryant, D. (2003). A classification of consensus methods for phylogenetics. In *Bioconsensus* (eds Janowitz, M.F., Lapointe, F.J., McMorris, F.R., Mirkin, B. and Roberts, F.S.), pp. 163–184. DIMACS Series in Discrete Mathematics and Theoretical Computer Science, vol. 61, Providence, RI, American Mathematical Society.
- Buneman, P. (1971). The recovery of trees from measures of dissimilarity. In *Mathematics in the Archaeological and Historical Sciences* (eds Hodson, F.R., Kendall, D.G. and Tautu, P.), pp. 387–395. Edinburgh, Edinburgh University Press.
- Burnham, K.P. and Anderson, D.R. (1998). *Model Selection and Inference. A Practical Information-theoretic Approach*. New York, Springer.
- Cameron, H.D. (1987). The upside-down cladogram: Problems in manuscript affiliation. In *Biological Metaphor and Cladistic Classification: An Interdisciplinary Perspective* (eds H.M. Hoenigswald and L.F. Wiener), pp. 227–242. Philadelphia, PA, University of Pennsylvania Press.
- Camin, J.H. and Sokal, R.R. (1965). A method for deducing branching sequences in phylogeny. *Evolution* **19**: 311–326.
- Carillo, H. and Lipman, D. (1988). The multiple sequence alignment problem in biology. *SIAM J. Appl. Math.* **48**: 1073–1082.
- Carroll, L. (1872). *Through the Looking Glass*. London, Macmillan.
- Carpenter, J.M. (2003). On “Molecular phylogeny of Vespidae (Hymenoptera) and the evolution of sociality in wasps”. *Am. Museum Novitates* **3389**: 1–20.
- Carter, M., Hendy, M.D., Penny, D., Székely, L.A. and Wormald, N.C. (1990). On the distribution of lengths of evolutionary trees. *SIAM J. Discrete Math.* **3**: 38–47.
- Cartmill, M. (1981). Hypothesis testing and phylogenetic reconstruction. *Z. Zool. Syst. Evol.-forsch.* **19**: 73–96.
- Chang, B.S.W., Jönsson K., Kazmi, M.A., Donoghue, M.J. and Sakmar T.P. (2002). Recreating a functional ancestral archosaur visual pigment. *Mol. Biol. Evol.* **19**: 1483–1489.
- Chang, J. (1996). Full reconstruction of Markov models on evolutionary trees: Identifiability and consistency. *Math. Biosc.* **137**: 51–73.
- Chang, J.T. and Kim, J. (1996). The measurement of homoplasy: A stochastic view. In *Homoplasy: the Recurrence of Similarity in Evolution* (eds M.J. Sanderson and L. Hufford), pp. 189–303. Academic Press.
- Chase, M.W., Soltis, D.E., Olmstead, R.G., Morgan, D., Les, D.H., Mishler, B.D., Duvall, M.R., Price, R.A., Hills, H.G., Qiu, Y.-L. *et al.* (1993). Phylogenetics of seed plants: An analysis of nucleotide sequences from the plastid gene *rbcL*. *Ann. Missouri Bot. Gard.* **40**: 528–580.

- Crichton, M. (1990). *Jurassic Park*. New York, Ballantine Books.
- Crick, F. (1968). The origin of the genetic code. *J. Mol. Biol.* **38**: 367–379.
- Crisci, J.V. and Stuessy, T.F. (1980). Determining primitive character states for phylogenetic reconstruction. *Syst. Bot.* **5**: 112–135.
- Cummings, M., Handley, S., Myers, D., Reed, D., Rokas, A. and Winka, K. (2003). Comparing bootstrap and posterior probability values in the four-taxon case. *Syst. Biol.* **52**: 477–487.
- Dacks, J.B. and Doolittle, W.F. (2001). Reconstructing/deconstructing the earliest eukaryotes: How comparative genomics can help. *Cell* **107**: 419–425.
- Dandekar, T., Snel, B., Huynen, M. and Bork, P. (1998). Conservation of gene order: A fingerprint of proteins that physically interact. *Trends Biochem. Sci.* **23**: 324–328.
- Darwin, C. (1859). *The Origin of Species by Means of Natural Selection, or the Preservation of Favoured Races in the Struggle for Life*. London, John Murray [1964. Facsimile of 1st edition. Cambridge, MA, Harvard University Press]
- Davids, W., Gamielien, J., Liberles, D.A. and Hide, W. (2002). Positive selection scanning reveals decoupling of enzymatic activities of carbamoyl phosphate synthetase in *H. pylori*. *J. Mol. Evol.* **54**: 458–464.
- Davidson, D. (1991). On the individuation of events. *Synthese* **86**: 229–254.
- Davis, J.I. and Nixon, K.C. (1992). Populations, genetic variation, and the delimitation of phylogenetic species. *Syst. Biol.* **41**: 421–435.
- Davis, J.I., Stevenson, D.W., Petersen, G., Seberg, O., Campbell, L.M., Freudenstein, J.V., Goldman, D.H., Hardy, C.R., Michelangeli, F.A., Simmons, M.P. *et al.* (2004). A phylogeny of the monocots, as inferred from *rbcL* and *atpA* sequence variation, and a comparison of methods for calculating jackknife and bootstrap values. *Syst. Bot.* **29**: 467–510.
- Dayhoff, M.O. and Eck, R.V. (1968). *Atlas of Protein Sequence and Structure*. 1967–68. Silver Spring, MD, National Biomed. Res. Foundation.
- Dayhoff, M.O. and Park, C.M. (1969). Cytochrome C: Building a phylogenetic hypothesis. In *Atlas of Protein Sequence and Structure*. 1969 (ed. M.O. Dayhoff), pp. 7–16 vol. 4. Silver Spring, MD, National. Biomed. Res. Foundation.
- DeBry, R.W. and Slade, N.A. (1985). Cladistic analysis of restriction endonuclease cleavage maps within a maximum-likelihood framework. *Syst. Zool.* **34**: 21–34.
- De Laet, J. (1997). *A Reconsideration of Three-Item Analysis, the Use of Implied Weights in Cladistics, and a Practical Application in Gentianaceae*. PhD Thesis, University of Leuven, Belgium.
- De Laet, J. (2003). Parsimony algorithms for characters that are inapplicable in some terminals (Abstract, 21st annual meeting of the Willi Hennig Society, Helsinki 2002). *Cladistics* **19**: 151.
- De Laet, J. (2004). When one and one is not two: Parsimony analysis of sequence data (Abstract, 22nd annual meeting of the Willi Hennig Society, New York 2003). *Cladistics* **20**: 81.
- De Laet, J. and Smets, E. (1998). On the three-taxon approach to parsimony analysis. *Cladistics* **14**: 363–381.
- De Laet, J. and Wheeler, W. (2003). *POY version 3.0.11* (Wheeler, Gladstein and De Laet, May 6 2003). Command line documentation. Available at <ftp://ftp.amnh.org/pub/molecular/poy>.
- de Pinna, M. C. C. (1991). Concepts and tests of homology in the cladistic paradigm. *Cladistics* **7**: 367–394.
- de Queiroz, K. (1992). Phylogenetic definitions and taxonomic philosophy. *Biol. Philos.* **7**: 295–313.
- de Queiroz, K. (1996). Including the characters of interest during tree reconstruction and the problems of circularity and bias in studies of character evolution. *Am. Nat.* **148**: 700–708.
- de Queiroz, K. and Poe, S. (2001). Philosophy and phylogenetic inference: A comparison of likelihood and parsimony methods in the context of Karl Popper's writings on corroboration. *Syst. Biol.* **50**: 305–321.
- de Queiroz, K. and Poe, S. (2003). Failed refutations: Further comments on parsimony and likelihood methods and their relationship to Popper's degree of corroboration. *Syst. Biol.* **52**: 352–367.
- Dezulian, T. and Steel, M. (2004). Phylogenetic closure operations and homoplasy-free evolution. In *Classification, Clustering, and Data Mining Applications* (Proceedings of the meeting of the International Federation of Classification Societies (IFCS) 2004). (eds D. Banks, L. House, F.R. McMorris, P. Arabie and W. Gaul), pp. 395–416. Springer-Verlag, Berlin.
- Dibb, N.J. and Newman, A.J. (1989). Evidence that introns arose at proto-splice sites. *EMBO J.* **8**: 2015–2021.
- Dollo, L. (1893). Le lois de l'évolution. *Bull. Soc. Belge Geol. Paleontol. d'Hydrog.* **7**: 164–167.
- Donoghue, M.J. and Doyle, J.A. (1989). Phylogenetic analysis of angiosperms and the relationships of Hamamelidae. In *Evolution, Systematics and Fossil History of the Hamamelidae*, vol.1 (eds P. Crane and S. Blackmore), pp. 17–45. Oxford, Clarendon Press.
- Donoghue, M.J. and Sanderson, M.J. (1992). The suitability of molecular and morphological evidence in reconstructing plant phylogeny. In *Molecular Systematics of Plants*. (eds P.S. Soltis, D.E. Soltis and J.J. Doyle), pp. 340–368, New York, Chapman & Hall.

- Donoghue, M.J., Doyle, J.A., Gauthier, J.A., Kluge, A.G. and Rowe, T. (1989). The importance of fossils in phylogeny reconstruction. *Annu. Rev. Ecol. Syst.* **20**: 431–460.
- Doolittle, W.F. (1999). Lateral genomics. *Trends Cell. Biol.* **9**: M5–M8.
- Doolittle, W.F., Boucher, Y., Nesbo, C.L., Douady, C.J., Andersson, J.O. and Roger, A.J. (2003). How big is the iceberg of which organellar genes in nuclear genomes are but the tip? *Phil. Trans. R. Soc. Lond. B Biol. Sci.* **358**: 39–57.
- Duret, L., Mouchiroud, D. and Gouy, M. (1994). HOVERGEN: A database of homologous vertebrate genes. *Nucleic Acids Res.* **22**: 2360–2365.
- Edwards, A.W.F. (1972). *Likelihood*. Cambridge, Cambridge University Press.
- Edwards, A.W.F. and Cavalli-Sforza, L.L. (1963). The reconstruction of evolution. *Heredity* **18**: 553.
- Edwards, A.W.F. and Cavalli-Sforza, L.L. (1964). Reconstruction of evolutionary trees. In *Phenetic and Phylogenetic Classification* (eds V.H. Heywood and J. McNeill), pp. 67–76 no. 6. London, Syst. Assoc. Publ.
- Eisen, M.B., Spellman, P.T., Brown, P.O. and Botstein, D. (1998). Cluster analysis and display of genome-wide expression patterns. *Proc. Natl. Acad. Sci. USA* **95**: 14863–14868.
- Enard, W., Khaitovich, P., Klose, J., Zöllner, S., Heissig, F., Giavalisco, P., Nieselt-Struwe, K., Muchmore, E., Varki, A., Ravid, R. *et al.* (2002). Intra and interspecific variation in primate gene expression patterns. *Science* **296**: 340–343.
- Endo, T., Ikee, K. and Gojobori, T. (1996). Large-scale search for genes on which positive selection may operate. *Mol. Biol. Evol.* **13**: 685–690.
- Endress, P.K. (1994). *Diversity and Evolutionary Biology of Tropical Flowers*. Cambridge, Cambridge University Press.
- Erdős, P.L. and Székely, L.A. (1992). Evolutionary trees: An integer multicommodity maxflow–min-cut theorem. *Adv. Appl. Math.* **13**: 375–389.
- Erdős, P.L. and Székely, L.A. (1993). Counting bichromatic evolutionary trees. *Discrete Appl. Math.* **47**: 1–8.
- Erdős, P.L., Steel, M.A., Székely, L.A. and Warnow, T. (1999). A few logs suffice to build (almost) all trees (Part 1). *Random Struct Algorithms* **14**: 153–184.
- Excoffier, L. and Smouse, P.E. (1994). Using allele frequencies and geographic subdivision to reconstruct gene trees within a species: Molecular variance parsimony. *Genetics* **136**: 343–359.
- Farris, J.S. (1966). Estimation of conservatism of characters by constancy within biological populations. *Evolution* **20**: 319–334.
- Farris, J.S. (1967). The meaning of relationship and taxonomic procedure. *Syst. Zool.* **16**: 44–51.
- Farris, J.S. (1969). A successive approximations approach to character weighting. *Syst. Zool.* **18**: 374–385.
- Farris, J.S. (1970). Methods for computing Wagner trees. *Syst. Zool.* **19**: 83–92.
- Farris, J.S. (1972). Estimating phylogenetic trees from distance matrices. *Am. Nat.* **106**: 645–668.
- Farris, J.S. (1973a). On the use of the parsimony criterion for inferring phylogenetic trees. *Syst. Zool.* **22**: 250–256.
- Farris, J.S. (1973b). A probability model for inferring evolutionary trees. *Syst. Zool.* **22**: 250–256.
- Farris, J.S. (1977a). Phylogenetic analysis under Dollo's law. *Syst. Zool.* **26**: 77–88.
- Farris, J.S. (1977b). Some further comments on Le Quesne's methods. *Syst. Zool.* **26**: 220–223.
- Farris, J.S. (1978a). Inferring phylogenetic trees from chromosome inversion data. *Syst. Zool.* **27**: 275–284.
- Farris, J.S. (1978b). *Wagner78*. Published by the author.
- Farris, J.S. (1979). The information content of the phylogenetic system. *Syst. Zool.* **28**: 483–519.
- Farris, J.S. (1982a). Outgroups and parsimony. *Syst. Zool.* **31**: 328–334.
- Farris, J.S. (1982b). Simplicity and informativeness in systematics and phylogeny. *Syst. Zool.* **31**: 413–444.
- Farris, J.S. (1983). The logical basis of phylogenetic analysis. In *Advances in Cladistics*, volume 2: *Proceedings of the Second Meeting of the Willi Hennig Society*. (eds N. Platnick and V. Funk) pp. 7–36. New York, Columbia University Press.
- Farris, J.S. (1983/1994). The logical basis of phylogenetic analysis. In *Advances in Cladistics — Proceedings of the 2nd Annual Meeting of the Willi Hennig Society* (eds N. Platnick and V. Funk. New York, Columbia University Press. Abridged and reprinted in E. Sober (ed.) pp. 7–36. *Conceptual Issues in Evolutionary Biology*, Cambridge, MA, MIT Press, 1994 (page references to the latter).
- Farris, J.S. (1986). On the boundaries of phylogenetic systematics. *Cladistics* **2**: 14–27.
- Farris, J.S. (1988). *Hennig86*. Published by the author, Port Jefferson Station, New York.
- Farris, J.S. (1989a). The retention index and the rescaled consistency index. *Cladistics* **5**: 417–419.
- Farris, J.S. (1989b). Entropy and fruit flies. *Cladistics* **5**: 103–108.
- Farris, J.S. (1991). Hennig defined paraphyly. *Cladistics* **7**: 297–304.
- Farris, J.S. (1997). Cycles. *Cladistics* **13**: 131–144.
- Farris, J.S. (1999). Likelihood and inconsistency. *Cladistics* **15**: 199–204.
- Farris, J.S. (2001). Support weighting. *Cladistics* **17**: 389–394.

- Farris, J.S. and Kluge, A.G. (1986). Synapomorphy, parsimony, and evidence. *Taxon* **35**: 298–315.
- Farris, J.S., Kluge, A.G. and Eckardt, M.J. (1970). A numerical approach to phylogenetic systematics. *Syst. Zool.* **19**: 172–189.
- Farris, J.S., Källersjö, M., Albert, V.A., Allard, M., Anderberg, A., Bowditch, B., Bult, C., Carpenter, J.M., Crowe, T.M., De Laet, J. *et al.* (1995). Explanation. *Cladistics* **11**: 211–218.
- Farris, J.S., Albert, V.A., Källersjö, M., Lipscomb, D. and Kluge, A.G. (1996). Parsimony jackknifing outperforms neighbor-joining. *Cladistics* **12**: 99–124.
- Farris, J.S., Källersjö, M. and De Laet, J.E. (2001a). Branch lengths do not indicate support — even in maximum likelihood. *Cladistics* **17**: 298–299.
- Farris, J.S., Kluge, A.G. and De Laet, J.E. (2001b). Taxic revisions. *Cladistics* **17**: 79–103.
- Fedorov, A., Merican, A.F. and Gilbert, W. (2002). Large-scale comparison of intron positions among animal, plant, and fungal genes. *Proc. Natl. Acad. Sci. USA* **99**: 16128–16133.
- Felsenstein, J. (1968). *Statistical Inference and the Estimation of Phylogenies*. PhD Thesis, University of Chicago, Chicago, IL.
- Felsenstein, J. (1973). Maximum likelihood and minimum-steps methods for estimating evolutionary trees from data on discrete characters. *Syst. Zool.* **22**: 240–249.
- Felsenstein, J. (1978a). Cases in which parsimony and compatibility methods can be positively misleading. *Syst. Zool.* **27**: 401–410.
- Felsenstein, J. (1978b). The number of evolutionary trees. *Syst. Zool.* **27**: 27–33.
- Felsenstein, J. (1979). Alternative methods of phylogenetic inference and their interrelationship. *Syst. Zool.* **28**: 49–62.
- Felsenstein, J. (1981a). Evolutionary trees from DNA sequences: A maximum likelihood approach. *J. Mol. Evol.* **17**: 368–376.
- Felsenstein, J. (1981b). Evolutionary trees from gene frequencies and quantitative characters: Finding maximum likelihood estimates. *Evolution* **35**: 1229–1242.
- Felsenstein, J. (1981c). A likelihood approach to character weighting and what it tells us about parsimony and compatibility. *Biol. J. Linn. Soc.* **16**: 183–196.
- Felsenstein, J. (1982). Numerical methods for inferring evolutionary trees. *Q. Rev. Biol.* **57**: 379–404.
- Felsenstein, J. (1983). Methods for inferring phylogenies: A statistical view. In *Numerical Taxonomy* (ed. J. Felsenstein), pp. 315–334. Berlin, Springer-Verlag.
- Felsenstein, J. (1988). Phylogenies from molecular sequences: Inference and reliability. *Annu. Rev. Genet.* **2**: 521–565.
- Felsenstein, J. (1996). Inferring phylogenies from protein sequences by parsimony, distance, and likelihood methods. *Methods Enzymol.* **266**: 418–427.
- Felsenstein, J. (2004). *Inferring Phylogenies*. Sunderland, MA, Sinauer Associates.
- Felsenstein, J. and Sober, E. (1987). Parsimony and likelihood: An exchange. *Syst. Zool.* **35**: 617–626.
- Feng, D.F. and Doolittle, R.F. (1987). Progressive sequence alignment as a prerequisite to correct phylogenetic trees. *J. Mol. Evol.* **25**: 351–360.
- Fink, W.L. (1982). The conceptual relationship between ontogeny and phylogeny. *Paleobiology* **8**: 254–264.
- Fitch, W.M. (1970). Distinguishing homologous from analogous proteins. *Syst. Zool.* **19**: 99–113.
- Fitch, W.M. (1971). Toward defining the course of evolution: Minimal change for a specific tree topology. *Syst. Zool.* **20**: 406–416.
- Fitz-Gibbon, S.T. and House, C.H. (1999). Whole genome-based phylogenetic analysis of free-living microorganisms. *Nucleic Acids Res.* **27**: 4218–4222.
- Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y.L. and Postlethwait, J. (1999). Preservation of duplicate genes by complementary, degenerative mutations. *Genetics* **151**: 1531–1545.
- Forster, M.R. (2000). Key concepts in model selection: Performance and generalizability. *J. Math. Psych.* **44**: 205–231.
- Forster, M.R. and Sober, E. (1994). How to tell when simpler, more unified, or less *ad hoc* theories will provide more accurate predictions. *Br. J. Phil. Sci.* **45**: 1–35.
- Foulds, L.R. (1984). Maximum savings in the Steiner problem in phylogeny. *J. Theoret. Biol.* **107**: 471–474.
- Foulds, L.R. and Graham, R.L. (1982). The Steiner problem in phylogeny is NP-complete. *Adv. Appl. Math.* **3**: 43–49.
- Friedman, M. (1983). *Foundations of Space-Time Theories: Relativistic Physics and Philosophy of Science*. Princeton, NJ, Princeton University Press.
- Fredman, M.L. (1984). Algorithms for computing evolutionary similarity measures with length independent gap penalties. *Bull. Math. Biol.* **46**: 545–563.
- Freudenstein, J.V., Pickett, K.M., Simmons, M.P. and Wenzel, J.W. (2003). From basepairs to birdsongs: phylogenetic data in the age of genomics. *Cladistics* **19**: 333–347.
- Frost, D.R. (2000). Species, descriptive efficiency, and progress in systematics. In *The Biology of Plethodontid Salamanders* (eds R.C. Bruce, R.J. Jaeger, and L.D. Houck), pp. 7–29. New York, Kluwer Academic/Plenum Publishing.

- Frost, D.R. and Kluge, A.G. (1994). A consideration of epistemology in systematic biology, with special reference to species. *Cladistics* **10**: 259–294.
- Frost, D.R., Rodrigues, M.T., Grant, T. and Titus, T.A. (2001). Phylogenetics of the lizard genus *Tropidurus* (Squamata: Tropiduridae: Tropidurinae): Direct optimization, descriptive efficiency, and sensitivity analysis of congruence between molecular data and morphology. *Mol. Phylogenet. Evol.* **21**: 352–371.
- Fukami-Kobayashi, K., Schreiber, D.R. and Benner, S.A. (2002). Detecting compensatory covariation signals in protein evolution using reconstructed ancestral sequences. *J. Mol. Biol.* **319**: 729–743.
- Funk, V.A. and Brooks, D.R. (1990). *Phylogenetic Systematics as the Basis of Comparative Biology*. Washington, DC, Smithsonian Institution Press.
- Gaasterland, T. and Ragan, M.A. (1998). Microbial genescapes: Phyletic and functional patterns of ORF distribution among prokaryotes. *Microb. Comp. Genomics* **3**: 199–217.
- Gallut, C. and Barriel, V. (2002). Cladistic coding of genomic maps. *Cladistics* **18**: 526–536.
- Galperin, M.Y. and Koonin, E.V. (2000). Who's your neighbor? New computational approaches for functional genomics. *Nat. Biotechnol.* **18**: 609–613.
- Gee, H. (2000). *Deep Time: Cladistics, the Revolution in Evolution*. London, Fourth Estate.
- Ghiselin, M.T. (1966). On semantic pitfalls of biological adaptation. *Philos. Sci.* **33**: 147–153.
- Ghiselin, M.T. (2004). Mayr and Bock versus Darwin on genealogical classification. *J. Zool. Syst. Evol. Res.* **42**: 165–169.
- Giribet, G. (2002). Relationships among metazoan phyla as inferred from 18S rRNA sequence data: A methodological approach. In *Molecular Systematics and Evolution: Theory and Practice*, (eds R. DeSalle, G. Giribet, and W. Wheeler), pp. 85–101. Basel, Birkhäuser Verlag.
- Giribet, G. and Wheeler, G. (1999). On gaps. *Mol. Phylogenet. Evol.* **13**: 132–143.
- Giribet, G., Distel, D.L., Polz, M., Sterrer, W. and Wheeler, W.C. (2000). Triploblastic relationships with emphasis on the acoelomates and the position of Gnathostomulida, Cyclophora, Plathelminthes, and Chaetognatha: A combined approach of 18S rDNA sequences and morphology. *Syst. Biol.* **49**: 539–562.
- Giribet, G., Wheeler, W.C. and Muona, J. (2002). DNA multiple sequence alignments. In *Molecular Systematics and Evolution: Theory and Practice* (eds R. Desalle, G. Giribet and W. Wheeler), pp. 107–114. Basel, Birkhäuser Verlag.
- Gladstein, D.S. (1997). Efficient incremental character optimization. *Cladistics* **13**: 21–26.
- Gogarten, J.P., Doolittle, W.F. and Lawrence, J.G. (2002). Prokaryotic evolution in light of gene transfer. *Mol. Biol. Evol.* **19**: 2226–2238.
- Goldman, N. (1990). Maximum likelihood inference of phylogenetic trees, with special reference to a Poisson process model of DNA substitution and to parsimony analyses. *Syst. Zool.* **39**: 345–361.
- Goloboff, P.A. (1995). A revision of the south American spiders of the family Nemesiidae (Araneae, Mygalomorphae). Part I: species from Peru, Chile, Argentina, and Uruguay. *Bull. Am. Mus. Nat. Hist.* **224**: 1–189.
- Goloboff, P.A. (1993a). Estimating character weights during tree search. *Cladistics* **9**: 83–91.
- Goloboff, P.A. (1993b). *Nona*: a tree-searching program. Available at <http://www.zmuc.dk/public/phylogeny/Nona-PeeWee/>.
- Goloboff, P.A. (1993c). *Pee-Wee*: Parsimony and Implied weights. Available at <http://www.zmuc.dk/public/phylogeny/Nona-PeeWee/>.
- Goloboff, P.A. (1994). Character optimization and calculation of tree lengths. *Cladistics* **9**: 433–436.
- Goloboff, P.A. (1995). *SPA: Sankoff Parsimony Analysis*, ver. 1.1. Available at <http://www.zmuc.dk/public/phylogeny/Nona-PeeWee/>.
- Goloboff, P.A. (1996a). *PHAST: Phylogenetic Analysis for Sankovian Transformations*, ver. 1.1. Available at <http://www.zmuc.dk/public/phylogeny/Nona-PeeWee/>.
- Goloboff, P.A. (1996b). Methods for faster parsimony analysis. *Cladistics* **12**: 199–220.
- Goloboff, P.A. (1998b). Tree searches under Sankoff parsimony. *Cladistics* **14**: 229–238.
- Goloboff, P.A. (1999). Analyzing large data sets in reasonable times: Solutions for composite optima. *Cladistics* **15**: 415–428.
- Goloboff, P.A. (2003). Parsimony, likelihood, and simplicity. *Cladistics* **19**: 91–103.
- Goloboff, P.A. and Farris, J.S. (2001). Methods for quick consensus estimation. *Cladistics* **17**: S26–S34.
- Goloboff, P.A., Wheeler, W. and Pol, D. (2003a). Parallel TNT. *Cladistics* **19**: 152 (in Muona, J. (2003). Abstracts of the 21st annual meeting of the Willi Hennig society. *Cladistics* **19**: 148–163.)
- Goloboff, P., Farris, J., Källersjö, M., Oxelmann, B., Ramírez, M. and Szumik, C. (2003b). Improvements to resampling measures of group support. *Cladistics* **19**: 324–332.
- Goloboff, P., Farris, J. and Nixon, K. (2004). *T.N.T.: Tree Analysis Using New Technology*. Available at www.zmuc.dk/public/phylogeny/tnt.
- Goodman, M., Czelusniak, J., Moore, G.W., Romero-Herrera, A.E. and Matsuda, G. (1979). Fitting the gene

- lineage into its species lineage, a parsimony strategy illustrated by cladograms constructed from globin sequences. *Syst. Zool.* **28**: 132–163.
- Gouldge, T.A. (1961). *The Ascent of Life. A Philosophical Study of the Theory of Evolution*. Toronto, University of Toronto Press. [1967 reprint]
- Grant, T. (2002). Testing methods: The evaluation of discovery operations in evolutionary biology. *Cladistics* **18**: 94–111.
- Grant, T. and Kluge, A.G. (2003). Data exploration in phylogenetic inference: Scientific, heuristic, or neither. *Cladistics* **19**: 379–418.
- Grant, T. and Kluge, A.G. (2004). Transformation series as an ideographic character concept. *Cladistics* **20**: 23–31.
- Greene, B. (2004). *The Fabric of the Cosmos. Space, Time, and the Texture of Reality*. New York, A.A. Knopf.
- Greuter, W., McNeill, J., Barrie, F.R., Burdet, H.M., Demoulin, V., Filgueiras, T.S., Nicolson, D.H., Silva, P.C., Skog, J.E., Trehane, P., et al. (2000). *International Code of Botanical Nomenclature (St. Louis Code)*. *Regnum Vegetabile* 138. Königstein, Koeltz Scientific Books.
- Gu, J. and Gu, X. (2003). Induced gene expression in human brain after the split from chimpanzee. *Trends Genet.* **19**: 63–65.
- Gusfield, D. (1997). *Algorithms on Strings, Trees, and Sequences: Computer Science and Computational Biology*. Cambridge, Cambridge University Press.
- Hacking, I. (1965). *The Logic of Statistical Inference*. Cambridge, Cambridge University Press.
- Hartigan, J.A. (1973). Minimum mutation fits to a given tree. *Biometrics* **29**: 53–65.
- Harvey, P.H. and Pagel, M.D. (1991). *The Comparative Method in Evolutionary Biology*. New York, Oxford University Press.
- Hasegawa, M. and Kishino, H. (1989). Confidence limits on the maximum-likelihood estimate of the hominoid tree from mitochondrial-DNA sequences. *Evolution* **43**: 672–677.
- Hastings, W.K. (1970). Monte Carlo sampling methods using Markov chains and their applications. *Biometrika* **57**: 97–109.
- Hedges, S.B. (2002). The origin and evolution of model organisms. *Nat. Rev. Genet.* **3**: 838–849.
- Hein, J. (1989a). A new method that simultaneously aligns and reconstructs ancestral sequences for any number of homologous sequences when a phylogeny is given. *Mol. Biol. Evol.* **6**: 649–668.
- Hein, J. (1989b). A tree reconstruction method that is economical in the number of pairwise comparisons used. *Mol. Biol. Evol.* **6**: 669–684.
- Hein, J.J. (2001). An algorithm for statistical alignment of sequences related by a binary tree. In *Pacific Symposium on Biocomputing 2001*. (eds R.B. Altman, A.K. Dunker, L. Hunter, K. Lauderdale and T.E. Klein), vol. 6, pp. 179–190. Singapore, World Scientific.
- Hein, J., Jensen, J.L. and Pedersen, C.N.S. (2003). Recursions for statistical multiple alignment. *Proc. Natl. Acad. Sci. USA* **100**: 14960–14965.
- Hendy, M.D. and Penny, D. (1982). Branch and bound algorithms to determine minimal evolutionary trees. *Math. Biosci.* **59**: 277–290.
- Hendy, M.D., Foulds, L.R. and Penny, D. (1980). Proving phylogenetic trees minimal with 1-clustering and set partitioning. *Math. Biosci.* **51**: 71–88.
- Hennig, W. (1950). *Grundzüge einer Theorie der phylogenetischen Systematik*. Berlin, Deutscher Zentralverlag.
- Hennig, W. (1966). *Phylogenetic Systematics*. Urbana, IL, University of Illinois Press.
- Higgins, D.G. and Sharp, P.M. (1988). CLUSTAL: A package for performing multiple sequence alignment on a microcomputer. *Gene* **73**: 237–244.
- Huber, K.T., Moulton, V. and Steel, M. (2002). *Four characters suffice to convexly define a phylogenetic tree*. Research Report UCDMA2002/12, Christchurch, New Zealand, Department of Mathematics and Statistics, University of Canterbury.
- Huelsenbeck, J.P. and Lander, K.M. (2003). Frequent inconsistency of parsimony under a simple model of cladogenesis. *Syst. Biol.* **52**: 641–648.
- Huelsenbeck, J.P. and Ronquist, F. (2001). MrBayes: Bayesian inference of phylogeny. *Bioinformatics* **17**: 754–755.
- Huelsenbeck, J.P., Bull, J.J. and Cunningham, C.W. (1996). Combining data in phylogenetic analysis. *Trends Ecol. Evol.* **4**: 152–158.
- Huelsenbeck, J., Ronquist, F., Nielsen, R. and Bollback, J. (2001). Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* **294**: 2310–2314.
- Huelsenbeck, J.P., Larget, B., Miller, R.E. and Ronquist, F. (2002). Potential applications and pitfalls of Bayesian inference of phylogeny. *Syst. Biol.* **51**: 673–688.
- Huelsenbeck, J., Larget, B. and Alfaro, M. (2004). Bayesian phylogenetic model selection using reversible jump Markov chain Monte Carlo. *Mol. Biol. Evol.* **21**: 1123–1133.
- Hull, D.L. (1967). Certainty and circularity in evolutionary taxonomy. *Evolution* **21**: 174–189.
- Hull, D.L. (1974). *Philosophy of Biological Science*. Englewood Cliffs, NJ, Prentice-Hall.
- Hull, D.L. (1975). Central subjects and historical narratives. *History and Theory: Studies Philos. Hist.* **14**: 253–274.
- Hull, D.L. (1977). The ontological status of species as evolutionary units. In *Foundational Problems in Special Sciences* (eds R. Butts and J. Hintikka), pp. 91–102. Dordrecht, D. Reidel Pub. Co.

- Hull, D.L. (1981). Historical narratives and integrating explanations. In *Pragmatism and Purpose: Essays Presented to Thomas A. Goudge* (eds L.W. Sumner, J.G. Slater and F. Wilson), pp. 172–188, 308–310. Toronto, University of Toronto Press.
- Hull, D.L. (1982). Exemplars and scientific change. *Proc. Biennial Mtg. Phil. Sci. Assoc.* **2**: 479–503.
- Hull, D.L. (1989). *The Metaphysics of Evolution*. Albany, NY, SUNY Press.
- Huson, D.H. and Steel, M. (2004). Phylogenetic trees based on gene content. *Bioinformatics* **20**: 2044–2049.
- Huynen, M.A. and Bork, P. (1998). Measuring genome evolution. *Proc. Natl. Acad. Sci. USA* **95**: 5849–5856.
- ICZN (1999). *International Code of Zoological Nomenclature, 4th Edn.* London, International Trust for Zoological Nomenclature.
- Jacob, F., Perrin, D., Sanchez, C. and Monod, J. (1960). L'Operon: groupe de genes a expression coordonnee par un operateur. *C. R. Seance Acad. Sci.* **250**: 1727–1729.
- Jenner, R.A. (2004). The scientific status of metazoan cladistics: why current reserach practice must change. *Zool. Scripta* **33**: 293–310.
- Jermann, T.M., Opitz, J.G., Stackhouse, J. and Benner, S.A. (1995). Reconstructing the evolutionary history of the artiodactyl ribonuclease superfamily. *Nature* **374**: 57–59.
- Jiang, T.L. and Lawler, E.L. (1994). Aligning sequences via an evolutionary tree: Computational complexity and approximation. In *Proceedings of the 26th ACM Symposium on the Theory of Computing*, pp. 760–769. New York, ACM.
- Källersjö, M., Farris, J.S., Chase, M.W., Bremer, B., Fay, M.F., Humphries, C.J., Petersen, G., Seberg, O. and Bremer, K. (1998). Simultaneous parsimony jackknife analysis of 2538 *rbcl* DNA sequences reveals support for major clades of green plants, land plants, seed plants and flowering plants. *Plant Syst. Evol.* **213**: 259–287.
- Källersjö, M., Albert, V.A. and Farris, J.S. (1999). Homoplasy increases phylogenetic structure. *Cladistics* **15**: 91–93.
- Kapitonov, V.V. and Jurka, J. (1999). The long terminal repeat of an endogenous retrovirus induces alternative splicing and encodes an additional carboxy-terminal sequence in the human leptin receptor. *J. Mol. Evol.* **48**: 248–251.
- Katinka, M.D., Duprat, S., Cornillot, E., Metenier, G., Thomarat, F., Prensier, G., Barbe, V., Peyretailade, E., Brottier, P., Wincker, P. *et al.* (2001). Genome sequence and gene compaction of the eukaryote parasite *Encephalitozoon cuniculi*. *Nature* **414**: 450–453.
- Kidd, K.K. and Sgaramella-Zonta, L.A. (1971). Phylogenetic analysis: Concepts and methods. *Am. J. Hum. Genet.* **23**: 235–252.
- Kim, J. (1996). General inconsistency conditions for maximum parsimony: Effects of branch lengths and increasing numbers of taxa. *Syst. Biol.* **45**: 363–374.
- Kimura, M. and Crow, J. (1964). The number of alleles that can be maintained in a finite population. *Genetics* **49**: 725–738.
- Kjer, K.M. (1995). Use of rRNA secondary structure in phylogenetic studies to identify homologous positions: An example of alignment and data presentation from the frogs. *Mol. Phylogenet. Evol.* **4**: 314–330.
- Kluge, A.G. (1988). Parsimony in vicariance biogeography: A quantitative method and a Greater Antillean example. *Syst. Zool.* **37**: 315–328.
- Kluge, A.G. (1989). A concern for evidence and a phylogenetic hypothesis of relationships among *Epicrates* (Boidae, Serpentes). *Syst. Zool.* **38**: 7–25.
- Kluge, A.G. (1997a). Testability and the refutation and corroboration of cladistic hypotheses. *Cladistics* **13**: 81–96.
- Kluge, A.G. (1997b). Sophisticated falsification and research cycles: Consequences for differential character weighting in phylogenetic systematics. *Zool. Scripta* **26**: 349–360.
- Kluge, A.G. (1999). The science of phylogenetic systematics: Explanation, prediction, and test. *Cladistics* **15**: 429–436.
- Kluge, A.G. (2001a). Parsimony with and without scientific justification. *Cladistics* **17**: 199–210.
- Kluge, A.G. (2001b). Philosophical conjectures and their refutation. *Syst. Biol.* **50**: 322–330.
- Kluge, A.G. (2002). Distinguishing “or” from “and” and the case for historical identification. *Cladistics* **18**: 585–593.
- Kluge, A.G. (2003a). On the deduction of species relationships: A précis. *Cladistics* **19**: 233–239.
- Kluge, A.G. (2003b). The repugnant and the mature in phylogenetic inference: A temporal similarity and historical identity. *Cladistics* **19**: 356–368.
- Kluge, A.G. (2004). On total evidence: For the record. *Cladistics* **20**: 205–207.
- Kluge, A.G. (2005). Taxonomy in theory and practice, with arguments for a new phylogenetic system of taxonomy. In *Ecology and Evolution in the Tropics: a Herpetological Perspective* (eds M.A. Donnelly, B.I. Crother, C. Guyer, M.H. Wake and M.E. White), pp. 7–47. Chicago, University of Chicago Press.
- Kluge, A.G. and Farris, J.S. (1969). Quantitative phyletics and the evolution of Anurans. *Syst. Zool.* **18**: 1–32.
- Kluge, A.G. and Farris, J.S. (1999). Taxic homology = overall similarity. *Cladistics* **15**: 205–212.
- Koonin, E.V., Aravind, L. and Kondrashov, A.S. (2000). The impact of comparative genomics on our understanding of evolution. *Cell* **101**: 573–576.

- Koonin, E.V., Makarova, K.S. and Aravind, L. (2001). Horizontal gene transfer in prokaryotes: Quantification and classification. *Annu. Rev. Microbiol.* **55**: 709–742.
- Koonin, E.V. and Galperin, M.Y. (2002). *Sequence-Evolution-Function. Computational Approaches in Comparative Genomics*. Kluwer Academic Publishers, New York.
- Koonin, E.V., Fedorova, N.D., Jackson, J.D., Jacobs, A.R., Krylov, D.M., Makarova, K.S., Mazumder, R., Mekhedov, S.L., Nikolskaya, A.N., Rao, B.S., *et al.* (2004). A comprehensive evolutionary classification of proteins encoded in complete eukaryotic genomes. *Genome Biol.* **5**: R7.
- Korbel, J.O., Snel, B., Huynen, M.A. and Bork, P. (2002). SHOT: A web server for the construction of genome phylogenies. *Trends Genet.* **18**: 158–162.
- Koshi, J.M. and Goldstein, R.A. (1996). Probabilistic reconstruction of ancestral protein sequences. *J. Mol. Evol.* **42**: 313–320.
- Kruskal, J. (1983). An overview of sequence comparison. In *Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison* (eds D. Sankoff and J. Kruskal), pp. 1–44. Stanford, CA, CSLI Publications (1999 reprint).
- Kumar, S., Tamura, K. and Nei, M. (1993). MEGA: *Molecular Evolutionary Genetics Analysis*, vers. 1.01. University Park, PA, Pennsylvania State University.
- Kunin, V. and Ouzounis, C.A. (2003). The balance of driving forces during genome evolution in prokaryotes. *Genome Res.* **13**: 1589–1594.
- Lande, R. (1976). Natural selection and random genetic drift in phenotypic evolution. *Evolution* **30**: 314–334.
- Larget, B. and Simon, D. (1999). Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. *Mol. Biol. Evol.* **16**: 750–759.
- Larson, A. and Losos, J.B. (1996). Phylogenetic systematics of adaptation. In *Adaptation* (eds M.R. Rose and G.V. Lauder), pp. 187–220. San Diego, CA, Academic Press.
- Laudan, L. (1990). *Science and Relativism: Some Key Controversies in the Philosophy of Science*. Chicago, University of Chicago Press.
- Laudan, R. (1990). What's so special about the past? In *Evolutionary Innovations* (ed. M. Nitechi), pp. 55–67. Chicago, University of Chicago Press.
- Lauder, G.V., Leroi, A.M. and Rose, M.R. (1993). Adaptations and History. *Trends Ecol. Evol.* **8**: 294–297.
- Lawrence, J. (1999). Selfish operons: The evolutionary impact of gene clustering in prokaryotes and eukaryotes. *Curr. Opin. Genet. Dev.* **9**: 642–648.
- Lawrence, J.G. and Roth, J.R. (1996). Selfish operons: Horizontal transfer may drive the evolution of gene clusters. *Genetics* **143**: 1843–1860.
- Le Cam, L. (1960). An approximation theorem for the Poisson binomial distribution. *Pacific J. Math.* **10**: 1181–1197.
- Le Quesne, W. (1969). A method of selection of characters in numerical taxonomy. *Syst. Zool.* **18**: 201–205.
- Lee, D.C. and Bryant, H.N. (1999). A reconsideration of the coding of inapplicable characters: Assumptions and problems. *Cladistics* **15**: 373–378.
- Levesque, M., Shasha, D., Kim, W., Surette, M.G. and Benfey, P.N. (2003). Trait-to-gene: a computational method for predicting the function of uncharacterized genes. *Curr. Biol.* **13**: 129–133.
- Lewis, P.O. (2001). A likelihood approach to estimating phylogeny from discrete morphological character data. *Syst. Biol.* **50**: 913–925.
- Li, S. (1996). *Phylogenetic Tree Construction using Markov Chain Monte Carlo*. PhD Dissertation, Ohio State University, Columbus, OH.
- Li, S., Pearl, D.K. and Doss, H. (2000). Phylogenetic tree construction using Markov chain Monte Carlo. *J. Am. Stat. Assoc.* **2000**: 493–508.
- Liberles, D.A. (2001). Evaluation of methods for determination of a reconstructed history of gene sequence evolution. *Mol. Biol. Evol.* **18**: 2040–2047.
- Liberles, D.A., Schreiber, D.R., Govindarajan, S., Chamberlin, S.G. and Benner, S.A. (2001). The Adaptive Evolution Database (TAED). *Genome Biol.* **2**(8): research0028.1– research0028.6.
- Liberles, D.A., Thoren, A., von Heijne, G. and Elofsson, A. (2002). The use of phylogenetic profiles for gene predictions. *Curr. Genomics* **3**: 131–137.
- Lidén, M. (1990). Replicators, hierarchy, and the species problem. *Cladistics* **6**: 183–186.
- Lipscomb, D.L. (1992). Parsimony, homology, and the analysis of multistate characters. *Cladistics* **8**: 45–65.
- Logsdon, Jr., J.M., Stoltzfus, A. and Doolittle, W.F. (1998). Molecular evolution: Recent cases of spliceosomal intron gain? *Curr. Biol.* **8**: R560–R63.
- Logsdon, Jr., J.M., Tyshenko, M.G., Dixon, C., J, D.J., Walker, V.K. and Palmer, J.D. (1995). Seven newly discovered intron positions in the triose-phosphate isomerase gene: Evidence for the introns-late theory. *Proc. Natl. Acad. Sci. USA* **92**: 8507–8511.
- Long, M. and Rosenberg, C. (2000). Testing the “proto-splice sites” model of intron origin: Evidence from analysis of intron phase correlations. *Mol. Biol. Evol.* **17**: 1789–1796.
- Lutzoni, F., Wagner, P., Reeb, V. and Zoller, S. (2000). Integrating ambiguously aligned regions of DNA sequences in phylogenetic analyses without violating positional homology. *Syst. Biol.* **49**: 628–651.
- Lynch, M. and Richardson, A.O. (2002). The evolution of spliceosomal introns. *Curr. Opin. Genet. Dev.* **12**: 701–710.

- Maddison, D.R. (1991). The discovery and importance of multiple islands of most-parsimonious trees. *Syst. Zool.* **40**: 315–328.
- Maddison, D.R. and Maddison, W.P. (2001). *MacClade 4: Analysis of Phylogeny and Character Evolution* (incl. vers. 4.03). Sunderland, MA, Sinauer Associates.
- Maddison, D.R., Swofford, D.L. and Maddison, W.P. (1997). NEXUS: An extensible file format for systematic information. *Syst. Biol.* **46**: 590–621.
- Maddison, W.P. (1991). Squared-change parsimony reconstructions of ancestral states for continuous-valued characters on a phylogenetic tree. *Syst. Zool.* **40**: 304–314.
- Maddison, W.P. (1993). Missing data versus missing characters in phylogenetic analysis. *Syst. Biol.* **42**: 576–581.
- Maddison, W.P. and Maddison, D.R. (1992). *MacClade 3: Analysis of Phylogeny and Character Evolution* (incl. vers. 3.04). Sunderland, MA, Sinauer Associates.
- Mallatt, J. and Winchell, C.J. (2002). Testing the new animal phylogeny: First use of combined large-subunit and small-subunit rRNA gene sequences to classify the protostomes. *Mol. Biol. Evol.* **19**: 289–301.
- Marchionni, M. and Gilbert, W. (1986). The triosephosphate isomerase gene from maize: Introns antedate the plant-animal divergence. *Cell* **46**: 133–141.
- Marcotte, E.M., Pellegrini, M., Thompson, M.J., Yeates, T.O. and Eisenberg, D.A. (1999). A combined algorithm for genome-wide prediction of protein function. *Nature* **402**: 83–86.
- Martin C. and Paz-Ares J. (1997). MYB transcription factors in plants. *Trends Plant Sci.* **13**: 67–73.
- Mau, B. (1996). *Bayesian Phylogenetic Inference via Markov Chain Monte Carlo Methods*. PhD Dissertation, University of Wisconsin, Madison, WI.
- Mau, B. and Newton, M. (1997). Phylogenetic inference for binary data on dendrograms using Markov chain Monte Carlo. *J. Comput. Graph. Stat.* **6**: 122–131.
- Mau, B., Newton, M. andarget, B. (1999). Bayesian phylogenetic inference via Markov chain Monte Carlo methods. *Biometrics* **55**: 1–12.
- Mayr, E. and Bock, W.J. (2002). Classification and other ordering systems. *J. Zool. Syst. Evol. Res.* **40**: 169–194.
- McAllister, J.W. (1996). *Beauty and Revolution in Science*. Ithaca, NY, Cornell University Press.
- McAllister, J.W. (2000). Unification of theories. In *A Companion to the Philosophy of Science* (ed. W.H. Newton-Smith), pp. 537–539. Oxford, Blackwell Publishing.
- McDade, L.A. (1992). Hybrids and phylogenetic systematics II. The impact of hybrids on cladistic analysis. *Evolution* **46**: 1329–1346.
- Messier, W. and Stewart, C.B. (1997). Episodic adaptive evolution of primate lysozymes. *Nature* **385**: 151–154.
- Metropolis, N., Rosenbluth, A.W., Rosenbluth, M.N., Teller, A.H. and Teller, E. (1953). Equations of state calculations by fast computing machines. *J. Chem. Phys.* **21**: 1087–1091.
- Mickevich, M.F. and Farris, J.S. (1981). *PHYSYS: Phylogenetic Analysis System*. Published by the authors.
- Miklós, I., Lunter, A. and Holmes, I. (2004). A “long indel” model for evolutionary sequence alignment. *Mol. Biol. Evol.* **21**: 529–540.
- Mindell, D.P. and Thacker, C.E. (1996). Rates of molecular evolution: Phylogenetic issues and applications. *Annu. Rev. Ecol. Syst.* **27**: 279–303.
- Mirkin, B.G., Fenner, T.I., Galperin, M.Y. and Koonin, E.V. (2003). Algorithms for computing parsimonious evolutionary scenarios for genome evolution, the last universal common ancestor and dominance of horizontal gene transfer in the evolution of prokaryotes. *BMC Evol. Biol.* **3**: 2.
- Mishler, B.D. (1994). Cladistic analysis of molecular and morphological data. *Am. J. Phys. Anthropol.* **94**: 143–156.
- Mishler, B.D. (1999). Getting rid of species? In *Species: New Interdisciplinary Essays* (ed. R. Wilson), pp. 307–315. Cambridge, MA, MIT Press.
- Mishler, B.D. (2000). Deep phylogenetic relationships among “plants” and their implications for classification. *Taxon* **49**: 661–683.
- Mishler, B.D. and Brandon, R.N. (1987). Individuality, pluralism, and the phylogenetic species concept. *Biol. Phil.* **2**: 397–414.
- Mishler, B.D. and De Luna, E. (1991). The use of ontogenetic data in phylogenetic analyses of mosses. *Adv. Bryol.* **4**: 121–167.
- Mishler, B.D. and Theriot, E. (2000a). The phylogenetic species concept (*sensu* Mishler and Theriot): Monophyly, apomorphy, and phylogenetic species concepts. In *Species Concepts and Phylogenetic Theory: A Debate*. (eds Q.D. Wheeler and R. Meier), pp. 44–54. New York, Columbia University Press.
- Mishler, B.D. and Theriot, E.G. (2000b). A critique from the Mishler and Theriot phylogenetic species concept perspective: Monophyly, apomorphy, and phylogenetic species concepts. In *Species Concepts and Phylogenetic Theory: A Debate* (eds Q.D. Wheeler and R. Meier), pp. 133–145. New York, Columbia University Press.
- Mishler, B.D. and Theriot, E.G. (2000c). A defense of phylogenetic species concept (*sensu* Mishler and Theriot): Monophyly, apomorphy, and phylogenetic species concepts. In *Species Concepts and Phylogenetic Theory: A Debate* (eds Q.D. Wheeler and R. Meier), pp. 179–184. New York, Columbia University Press.
- Modrek, B. and Lee, C.J. (2003). Alternative splicing in the human, mouse, and rat genomes is associated with

- an increased frequency of exon creation and/or loss. *Nat. Genet.* **34**: 177–180.
- Moilanen, A. (1999). Searching for most parsimonious trees with simulated evolutionary optimization. *Cladistics* **15**: 39–50.
- Moilanen, A. (2001). Simulated evolutionary optimization and local search: Introduction and application to tree search. *Cladistics* **17**: S12–S25.
- Montague, M.G. and Hutchison, 3rd., C.A. (2000). Gene content phylogeny of herpesviruses. *Proc. Natl. Acad. Sci. USA* **97**: 5334–5339.
- Moran, N.A. (2002). Microbial minimalism: Genome reduction in bacterial pathogens. *Cell* **108**: 583–586.
- Moret, B.M.E., Wang, L.S., Warnow, T. and Wyman, S. (2001). New approaches for reconstructing phylogenies based on gene order. Proc. 9th Int'l Conf. on Intelligent Systems for Molecular Biology ISMB-2001, *Bioinformatics* **17**: S165–S173.
- Moret, B.M.E., Tang, J., Wang, L.S. and Warnow, T. (2002). Steps toward accurate reconstruction of phylogenies from gene-order data. *J. Comput. Syst. Sci.* **65**(3): 508–525.
- Morgenstern, B. (2004). DIALIGN: Multiple DNA and protein sequence alignment at BiBiServ. *Nucleic Acids Res.* **32**: W33–W36.
- Morgenstern, B., Dress, A. and Werner, T. (1996). Multiple DNA and protein sequence alignment based on segment-to-segment comparison. *Proc. Natl. Acad. Sci. USA* **93**: 12098–12103.
- Moritz, C. (2002). Strategies to protect biological diversity and the processes that sustain it. *Syst. Biol.* **51**: 238–254.
- Mossel, E. and Steel, M. (2004a). A phase transition for a random cluster model on phylogenetic trees. *Math. Biosci.* **187**: 189–203.
- Mossel, E. and Steel, M. (2005). How much can evolved characters tell us about the tree that generated them? In *Mathematics of Evolution and Phylogeny* (ed. O. Gascuel). Oxford, Oxford University Press.
- Murata, M., Richardson, J. and Sussman, J. (1985). Simultaneous comparison of three protein sequences. *Proc. Natl. Acad. Sci. USA* **82**: 3073–3077.
- Mushegian, A.R. and Koonin, E.V. (1996). Gene order is not conserved in bacterial evolution. *Trends Genet.* **12**: 289–290.
- Mushegian, A.R., Garey, J.R., Martin, J. and Liu, L.X. (1998). Large-scale taxonomic profiling of eukaryotic model organisms: A comparison of orthologous proteins encoded by the human, fly, nematode, and yeast genomes. *Genome Res.* **8**: 590–598.
- Myllykallio, H., Lipowski, G., Leduc, D., Filee, J., Forterre, P. and Liebl, U. (2002). An alternative flavin-dependent mechanism for thymidylate synthesis. *Science* **297**: 105–107.
- Nadeau, J.J. and Taylor, B.A. (1984). Lengths of chromosome segments conserved since divergence of man and mouse. *Proc. Natl. Acad. Sci. USA* **81**: 814–818.
- Naylor, G.J.P. and Adams D.C. (2001). Are the fossil data really at odds with the molecular data? Morphological evidence for *Cetartiodactyla* phylogeny reexamined. *Syst. Biol.* **50**: 444–453.
- Naylor, G.J.P. and Adams, D.C. (2003). Total evidence versus relevant evidence: A response to O'Leary *et al.* (2003). *Syst. Biol.* **52**: 864–865.
- Needleman, S.B. and Wunsch, C.D. (1970). A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Mol. Biol.* **48**: 443–453.
- Neff, N.A. (1986). A rational basis for *a priori* character weighting. *Syst. Zool.* **35**: 102–109.
- Nei, M. and Kumar, S. (2001). *Molecular Evolution and Phylogenetics*. Oxford, Oxford University press.
- Nelson, G. and Platnick, N.I. (1981). *Systematics and Biogeography: Cladistics and Vicariance*. New York, Columbia University Press.
- Newton, M., Mau, B. and Larget, B. (1999). Markov chain Monte Carlo for the Bayesian analysis of evolutionary trees from aligned molecular sequences. In *Statistics in Molecular Biology and Genetics*, vol. 33 (ed. F. Seillier-Moisewitsch), pp. 143–162. Bethesda, MD, Institute of Mathematical Statistics.
- Neyman, J. (1971). Molecular studies of evolution: A source of novel statistical problems. In *Statistical Decision Theory and Related Topics* (eds S. Gupta and J. Yackel), pp. 1–27. New York, Academic Press.
- Nixon, K.C. (1999). The parsimony ratchet, a new method for rapid parsimony analysis. *Cladistics* **15**: 407–414.
- Nixon, K.C. (2002). *WinClada*, vers. 1.00.08. Published by the author, Ithaca, New York (distributed through www.cladistics.org).
- Nixon, K.C. and Carpenter, J.M. (1993). On outgroups. *Cladistics* **9**: 413–426.
- Nixon, K.C. and Little, D.P. (2004). The use of optimality criteria in DNA sequence data and its application in a new computer program (Abstract, 22nd annual meeting of the Willi Hennig Society, New York 2003). *Cladistics* **20**: 90–91.
- Nixon, K.C. and Wheeler, Q.D. (1990). An amplification of the phylogenetic species concept. *Cladistics* **6**: 211–223.
- Nolan, D. (1997). Quantitative parsimony. *Br. J. Phil. Sci.* **48**: 329–343.

- Notredame, C. (2002). Recent progress in multiple sequence alignment: A survey. *Pharmacogenomics* **3**: 1–14.
- Notredame, C., Holm, L. and Higgins, D.G. (1998). COFFEE: An objective function for multiple sequence alignments. *Bioinformatics* **14**: 407–422.
- Notredame, C., Higgins, D.G. and Heringa, J. (2000). T-Coffee: A novel method for fast and accurate multiple sequence alignment. *J. Mol. Biol.* **302**: 205–217.
- O'Hara, R.J. (1988). Homage to Clío, or, toward an historical philosophy for evolutionary biology. *Syst. Zool.* **37**: 142–155.
- Ochman, H., Lawrence, J.G. and Groisman, E.A. (2000). Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**: 299–304.
- Ochoterena, H. (2004). Independence of alignment and phylogenetic reconstruction and their optimality criteria (Abstract, 22nd annual meeting of the Willi Hennig Society, New York 2003). *Cladistics* **20**: 91.
- Ohno, S. (1970). *Evolution by Gene Duplication*. New York, Springer-Verlag.
- Oleksiak, M.F., Churchill, G.A. and Crawford, D.L. (2002). Variation in gene expression within and among natural populations. *Nat. Genet.* **32**: 261–266.
- Padian, K. (2004). For Darwin, 'genealogy alone' did give classification. *J. Zool. Syst. Evol. Res.* **42**: 162–164.
- Parzen, E. (1962). *Stochastic Processes*. San Francisco, Holden-Day.
- Patterson, C. (1982). Morphological characters and homology. In *Problems of Phylogenetic Reconstruction*. (eds K.A. Joysey and A.E. Friday), pp. 21–74. New York, Academic Press.
- Patterson, C. (1988). The impact of evolutionary theories on systematics. In *Prospects in Systematics* (ed. D.L. Hawksworth), pp. 59–91. Syst. Assoc. Spec. Vol. 36. Oxford, Clarendon Press.
- Pellegrini, M., Marcotte, E.M., Thompson, M.J., Eisenberg, D. and Yeates, T.O. (1999). Assigning protein functions by comparative genome analysis: Protein phylogenetic profiles. *Proc. Natl. Acad. Sci. USA* **96**: 4285–4288.
- Penny, D., Lockhart, P.J., Steel, M.A. and Hendy, M.D. (1994). The role of models in reconstructing evolutionary trees. In *Models in Phylogeny Reconstruction* (eds R.W. Scotland, D.J. Siebert and D.M. Williams), pp. 211–230. Systematics Assoc. Special vol. 52. Oxford, Clarendon Press.
- Penny, D., Hendy, M.D., Lockhart, P.J. and Steel, M.A. (1996). Corrected parsimony, minimum evolution, and Hadamard conjugations. *Syst. Biol.* **45**: 596–606.
- Peterson, K.J. and Eernisse, D.J. (2001). Animal phylogeny and the ancestry of bilaterians: Inferences from morphology and 18S rDNA gene sequences. *Evol. Dev.* **3**: 170–205.
- Phillips, A., Janies, D. and Wheeler, W.C. (2000). Multiple sequence alignment in phylogenetic analysis. *Mol. Phylogenet. Evol.* **16**: 317–330.
- Planet, P.J., DeSalle, R., Siddall, M., Bael, T., Sarkar, I.N. and Stanley, S.E. (2001). Systematic analysis of DNA microarray data: Ordering and interpreting patterns of gene expression. *Genome Res.* **11**: 1149–1155.
- Platnick, N.I. (1979). Philosophy and the transformation of cladistics. *Syst. Zool.* **28**: 537–546.
- Platnick, N.I. and Cameron, H.D. (1977). Cladistic methods in textual, linguistic, and phylogenetic analysis. *Syst. Zool.* **26**: 380–385.
- Platnick, N.I., Griswold, C.E. and Coddington, J.A. (1991). On missing entries in cladistic analysis. *Cladistics* **7**: 337–343.
- Pleijel, F. (1995). On character coding for phylogeny reconstruction. *Cladistics* **11**: 309–315.
- Pol, D. and Siddall, M. (2001). Biases in maximum likelihood and parsimony: A simulation approach to a 10-taxon case. *Cladistics* **17**: 266–281.
- Popper, K. (1957). *The Poverty of Historicism*. London, Routledge and Kegan Paul.
- Popper, K. (1959). *The Logic of Scientific Discovery*. New York, Harper and Row [1968 edition].
- Popper, K. (1962a). Some comments on truth and the growth of knowledge. In *Logic, Methodology and Philosophy of Science* (eds E. Nagel, P. Suppes and A. Tarski), pp. 285–292. Proc 1960 Internatl. Congress. Stanford, CA, Stanford University Press.
- Popper, K. (1962b). *Conjectures and Refutations: The Growth of Scientific Knowledge*. London, Routledge and Kegan Paul.
- Popper, K. (1968). *The Logic of Scientific Discovery*. New York, Harper Torchbooks.
- Popper, K. (1979). *Objective Knowledge. An Evolutionary Approach*. New York, Oxford University Press.
- Popper, K. (1980). Evolution. *New Scientist* **87**: 611.
- Popper, K. (1983). *Realism and the Aim of Science*. London, Routledge.
- Posada, D. and Crandall, K. (1998). MODELTEST: Testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Posada, D. and Crandall, K. (2001a). Selecting models of nucleotide substitution: An application to human immunodeficiency virus 1 (HIV-1). *Mol. Biol. Evol.* **18**: 897–906.
- Pritchard, P.C.H. (1994). Cladistics: The great delusion. *Herpetol. Rev.* **25**: 103–110.
- Posada, D. and Crandall, K. (2001b). Selecting the best-fit model of nucleotide substitution. *Syst. Biol.* **50**: 580–601.
- Prömel, H.J. and Steger, A. (2000). A new approximation algorithm for the Steiner tree problem with performance ratio 5/3. *J. Algorithms* **36**: 89–101.

- Pupko, T., Pe'er, I., Shamir, R. and Graur, D. (2000). A fast algorithm for joint reconstruction of ancestral amino acid sequences. *Mol. Biol. Evol.* **17**: 890–896.
- Pupko, T., Pe'er, I., Hasegawa M., Graur, D. and Friedman, N. (2002). A branch-and-bound algorithm for the inference of ancestral amino-acid sequences when the replacement rate varies among sites: Application to the evolution of five gene families. *Bioinformatics* **18**: 1116–1123.
- Quine, W.V. (1963). On simple theories of a complex world. *Synthese* **15**: 103–106.
- Raff, R.A. (1996). *The Shape of Life: Genes, Development, and the Evolution of Animal Form*. Chicago, IL, University of Chicago Press.
- Rain, J.-C., Selig, L., De Reuse, H., Battaglia, V., Reverdy, C., Simon, S., Lenzen, G., Petel, F., Wojcik, J., Schachter, V. et al. (2001). The protein-protein interaction map of *Helicobacter pylori*. *Nature* **409**: 211–215.
- Rannala B. and Yang, Z. (1996). Probability distribution of molecular evolutionary trees: A new method of phylogenetic inference. *J. Mol. Evol.* **43**: 304–311.
- Reichenbach, H. (1956). *The Direction of Time*. Berkeley, CA, University of California Press.
- Remane, A. (1952). *Die Grundlagen des natürlichen Systems, der vergleichenden Anatomie und der Phylogenetik*. Leipzig, Akademische Verlagsgesellschaft Geest & Portig.
- Resch, A., Xing, Y., Alekseyenko, A., Modrek, B. and Lee, C. (2004). Evidence for a subpopulation of conserved alternative splicing events under selection pressure for protein reading frame conservation. *Nucleic Acids Res.* **32**: 1261–1269.
- Rexová, K., Frynta, D. and Zrzavý, J. (2003). Cladistic analysis of languages: Indo-European classification based on lexicostatistical data. *Cladistics* **19**: 120–127.
- Rice, K.A., Donoghue, M.J. and Olmstead, R.G. (1997). Analyzing large data sets: *rbcL* 500 revisited. *Syst. Biol.* **46**: 554–563.
- Rieppel, O.C. (1988). *Fundamentals of Comparative Biology*. Basel, Birkhäuser Verlag.
- Rieppel, O. (2003). Semaphoronts, cladograms and the roots of total evidence. *Biol. J. Linn. Soc.* **80**: 167–186.
- Rieppel, O. and Kearney, M. (2002). Similarity. *Biol. J. Linn. Soc.* **75**: 59–82.
- Rieseberg, L.H. and Soltis, D.E. (1991). Phylogenetic consequences of cytoplasmic gene flow in plants. *Evol. Trends Plants* **5**: 5–84.
- Rogers, J. (1997). On the consistency of the maximum likelihood estimation of phylogenetic trees from nucleotide sequences. *Syst. Biol.* **46**: 354–357.
- Rogozin, I.B., Wolf, Y.I., Sorokin, A.V., Mirkin, B.G. and Koonin, E.V. (2003). Remarkable interkingdom conservation of intron positions and massive, lineage-specific intron loss and gain in eukaryotic evolution. *Curr. Biol.* **13**: 1512–1517.
- Rokas, A. and Holland, P.W.H. (2000). Rare genomic changes as a tool for phylogenetics. *Trends Ecol. Evol.* **15**: 454–459.
- Romero, I., Fuertes, A., Benito, M.J., Malpica, J. M., Leyva, A. and Paz-Ares, J. (1998). More than 80 R2R3-MYB regulatory genes in the genome of *Arabidopsis thaliana*. *Plant J.* **14**: 273–284.
- Rossnes, R. (2004). *Ancestral Reconstruction of Continuous Characters and its Potential Application to Gene Expression and Alternative Splicing Analysis*. MSc Thesis, University of Bergen, Norway.
- Roth, V.L. (1984). On homology. *Biol. J. Linn. Soc.* **22**: 13–29.
- Roth, V.L. (1988). The biological basis of homology. In *Ontogeny and Systematics* (ed. Humpries, C.J.). New York, Columbia University Press.
- Royall, R. (1997). *Statistical Evidence — a Likelihood Paradigm*. New York, Chapman and Hall.
- Ruse, M. (1971). Narrative explanation and the theory of evolution. *Can. J. Phil.* **1**: 59–74.
- Russell, B. (1948). *Human Knowledge, its Scope and Limits*. London, George Allen and Unwin.
- Rutishauser, R. and Sattler, R. (1989). Complementary and heuristic value of contrasting models in structural biology. III. Case study on shoot-like “leaves” and leaf-like “shoots” in *Utricularia macrorhiza* and *Utricularia purpurea* (Lentibulariaceae). *Botanische Jahrbücher für Systematik* **111**: 121–137.
- Rzhetsky, A. and Nei, M. (1992). A simple method for estimating and testing minimum-evolution trees. *Mol. Biol. Evol.* **9**: 945–967.
- Rzhetsky, A., Ayala, F.J., Hsu, L.C., Chang, C. and Yoshida, A. (1997). Exon/intron structure of aldehyde dehydrogenase genes supports the “introns-late” theory. *Proc. Natl. Acad. Sci. USA* **94**: 6820–6825.
- Saitou, N. and Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406–425.
- Salem, A.H., Ray, D.A., Xing, J., Callinan, P.A., Myers, J.S., Hedges, D.J., Garber, R.K., Witherspoon, D.J., Jorde, L.B. and Batzer, M.A. (2003). Alu elements and hominid phylogenetics. *Proc. Natl. Acad. Sci. USA* **100**: 12787–12791.
- Salisbury, B.A. (1999). Strongest evidence: Maximum apparent phylogenetic signal as a new cladistic optimality criterion. *Cladistics* **15**: 137–149.
- Salmon, W.C. (1966). *The Foundations of Scientific Inference*. Pittsburgh, PA, University of Pittsburgh Press.
- Sanderson, M. and Hufford, L., eds (1996). *Homoplasy*. San Diego, CA, Academic Press.

- Sanderson, M. and Kim, J. (2000). Parametric phylogenetics? *Syst. Biol.* **49**: 817–829.
- Sanderson, M.J., Purvis, A. and Henze, C. (1998). Phylogenetic supertrees: Assembling the trees of life. *Trends Ecol. Evol.* **13**: 105–109.
- Sankoff, D. (1975). Minimal mutation trees of sequences. *SIAM J. Appl. Math.* **28**: 35–42.
- Sankoff, D. and Blanchette, M. (1998). Multiple genome rearrangement and breakpoint phylogeny. *J. Comp. Biol.* **5**: 555–570.
- Sankoff, D. and Cedergren, R.J. (1983). Simultaneous comparison of three or more sequences related by a tree. In *Time Warps, String Edits, and Macromolecules. The Theory and Practice of Sequence Comparison* (eds D. Sankoff, and J. Kruskal), pp. 253–263. Stanford, CA, CSLI Publications (1999 reprint).
- Sankoff, D. and Nadeau, J.H. (eds) (2000). *Comparative Genomics. Empirical and Analytical Approaches to Gene Order Dynamics, Map Alignment and the Evolution of Gene Families*. Dordrecht, Kluwer Academic Publishers.
- Sankoff, D. and Rousseau, P. (1975). Locating the vertices of a Steiner tree in arbitrary space. *Math. Program.* **9**: 240–246.
- Sankoff, D., Cedergren, R.J. and Lapalme, G. (1976). Frequency of insertion-deletion, transversion, and transition in the evolution of 5S ribosomal RNA. *J. Mol. Evol.* **7**: 133–149.
- Sankoff, D., Morel, C. and Cedergren, R.J. (1973). Evolution of 5S RNA and the non-randomness of base replacement. *Nat. New Biol.* **245**: 232–234.
- Sarkar, I.N., Planet, P.J., Bael, T.E., Stanley, S.E., Siddall, M., DeSalle, R. and Figurski, D.H. (2002). Characteristic attributes in cancer microarrays. *J. Biomed. Inform.* **35**: 111–122.
- Schwikowski, B. and Vingron, M. (1997). The deferred path heuristic for the generalized tree alignment problem. *J. Comput. Biol.* **4**: 415–431.
- Schwikowski, B. and Vingron, M. (2003). Weighted sequence graphs: Boosting iterated dynamic programming using locally suboptimal solutions. *Discr. Appl. Math.* **127**: 95–117.
- Scriven, M. (1959). Explanation and prediction in evolutionary theory. *Science* **130**: 477–482.
- Seitz, V., Ortiz García, S. and Liston, A. (2000). Alternative coding strategies and the inapplicable data coding problem. *Taxon* **49**: 47–54.
- Sellers, P.H. (1974). An algorithm for the distance between two sequences. *J. Comb. Theory* **16**: 253–258.
- Semple, C. and Steel, M. (2002). Tree reconstruction from multi-state characters. *Adv. Appl. Math.* **28**: 169–184.
- Semple, C. and Steel, M. (2003). *Phylogenetics*. Oxford, Oxford University Press.
- Shenkin, P.S., Erman, B. and Mastrandrea, L.D. (1991). Information-theoretical entropy as a measure of sequence variability. *Proteins Struct. Funct. Genet.* **11**: 297–313.
- Sicheritz-Ponten, T. and Andersson, S.G. (2001). A phylogenomic approach to microbial evolution. *Nucleic Acids Res.* **29**: 545–552.
- Siddall, M. (1998). Success of parsimony in the four-taxon case: Long branch repulsion by likelihood in the Farris zone. *Cladistics* **14**: 209–220.
- Siddall, M.E. and Kluge, A.G. (1997). Probabilism and phylogenetic inference. *Cladistics* **13**: 313–336.
- Sikes, D.S. and Lewis, P.O. (2001). *PAUPRat: PAUP Implementation of the Parsimony Ratchet*. Published by the authors (distributed through www.ucalgary.ca/~dsikes/sikes_lab.htm).
- Simmons, M.P. (2004). Independence of alignment and tree search. *Mol. Phylogenet. Evol.* **31**: 874–879.
- Simmons, M.P. and Ochoterena, H. (2000). Gaps as characters in sequence-based phylogenetic analyses. *Syst. Biol.* **49**: 369–381.
- Simon, D. and Larget, B. (1998). *Bayesian Analysis in Molecular Biology and Evolution (BAMBE)*, version 1.01 beta. Pittsburgh, PA, Department of Mathematics and Computer Science, Duquesne University.
- Simpson, G.G. (1964). *This View of Life: The World of an Evolutionist*. New York, Harcourt, Brace and World.
- Sinsheimer, J., Lake, J.A. and Little, R.J.A. (1996). Bayesian hypothesis testing of four-taxon topologies using molecular sequence data. *Biometrics* **52**: 193–210.
- Slatkin, M. and Maddison, W. (1989). A cladistic measure of gene flow inferred from the phylogenies of alleles. *Genetics* **123**: 603–613.
- Slowinski, J.B. (1998). The number of multiple alignments. *Mol. Phyl. Evol.* **10**: 264–266.
- Smith R.L. and Sytsma, K.J. (1990). Evolution of *Populus nigra* (sect. *Aigeiros*): Introgressive hybridization and the chloroplast contribution of *Populus alba* (sect. *Populus*). *Am. J. Bot.* **77**: 1176–1187.
- Smith, T.F., Waterman, M.S. and Fitch, W.M. (1981). Comparative biosequence metrics. *J. Mol. Evol.* **18**: 38–46.
- Smith, V.S., Page, R.D.M. and Johnson, K.P. (2004). Data incongruence and the problem of avian louse phylogeny. *Zool. Scripta* **33**: 239–259.
- Smouse, P.E. and Li, W.-H. (1989). Likelihood analysis of mitochondrial restriction-cleavage patterns for the human-chimpanzee-gorilla trichotomy. *Evolution* **41**: 1162–1176.
- Snel, B., Bork, P. and Huynen, M.A. (1999). Genome phylogeny based on gene content. *Nat. Genet.* **21**: 108–110.
- Snel, B., Bork, P. and Huynen, M.A. (2002). Genomes in flux: The evolution of archaeal and proteobacterial gene content. *Genome Res.* **12**: 17–25.

- Sober, E. (1980). Evolution, population thinking, and essentialism. *Phil. Sci.* **47**: 350–383.
- Sober, E. (1981). The principle of parsimony. *Br. J. Phil. Sci.* **32**: 145–156.
- Sober, E. (1983). Parsimony methods in systematics — philosophical issues. *Annu. Rev. Ecol. Syst.* **14**: 335–357.
- Sober, E. (1985). A likelihood justification for parsimony. *Cladistics* **1**: 209–233.
- Sober, E. (1986). Parsimony and character weighting. *Cladistics* **2**: 28–42.
- Sober, E. (1988a). *Reconstructing the Past: Parsimony, Evolution and Inference*. Cambridge, MA, MIT Press.
- Sober, E. (1988b). The conceptual relationship of cladistic phylogenetics and vicariance biogeography. *Syst. Zool.* **37**: 245–253.
- Sober, E. (1993). *Philosophy of Biology*. San Francisco, CA, Westview Press.
- Sober, E. (1994). Let's razor Ockham's Razor. In *Explanation and its Limits* (ed. D. Knowles), pp. 73–93. *Suppl. Philosophy, Roy. Inst. Phil.* **27**.
- Sober, E. (1996). Parsimony and predictive equivalence. *Erkenntnis* **44**: 167–197.
- Sober, E. (2002). Reconstructing ancestral character states — a likelihood perspective on cladistic parsimony. *The Monist* **85**: 156–176.
- Sober, E. (2003). Parsimony. In *The Philosophy of Science: an Encyclopedia* (eds S. Sarkar and J. Pfeifer). London, Routledge.
- Sober, E. (2004a). The contest between likelihood and parsimony. *Syst. Biol.* **53**: 644–653.
- Sober, E. (in press): Is drift a serious alternative to natural selection as an explanation of complex adaptive traits? In *The Spandrels of San Marco Twenty-Five Years After* (ed. D. Walsh). Oxford, Oxford University Press.
- Sober, E. and Steel, M. (2002). Testing the hypothesis of common ancestry. *J. Theor. Biol.* **218**: 395–408.
- Sokal, R.R. (1986). Phenetic taxonomy: Theory and methods. *Annu. Rev. Ecol. Syst.* **17**: 423–442.
- Sokal, R.R. and Camin, J.H. (1965). The two taxonomies: Areas of agreement and conflict. *Syst. Zool.* **14**: 176–195.
- Soltis, D.E., Soltis, P.S., Chase, M.W., Mort, M.E., Albach, D.C., Zanis, M., Savolainen, V., Hahn, W.H., Hoot, S.B., Fay, M.F. *et al.* (2000). Angiosperm phylogeny inferred from 18S rDNA, *rbcL*, and *atpB* sequences. *Bot. J. Linn. Soc.* **133**: 381–461.
- Sonnhammer, E.L. and Koonin, E.V. (2002). Orthology, paralogy and proposed classification for paralog subtypes. *Trends Genet.* **18**: 619–620.
- Steel, M.A. (1993). Distributions on bicoloured binary trees arising from the principle of parsimony. *Discrete Appl. Math.* **41**: 245–261.
- Steel, M. (2002). Some statistical aspects of the maximum parsimony method. In *Molecular Systematics and Evolution: Theory and Practice* (eds R. De Salle, R. Giribet and W. Wheeler), pp. 125–139. Basel, Birkhäuser Verlag.
- Steel, M. and Penny, D. (2000). Parsimony, likelihood, and the role of models in molecular phylogenetics. *Mol. Biol. Evol.* **17**: 839–850.
- Steel, M. and Penny, D. (2004). Two links between MP and ML under the Poisson model. *Applied Math. Lett.* (in press).
- Steel, M., Penny, D. and Hendy, M. (1993). Parsimony can be consistent! *Syst. Biol.* **42**: 581–587.
- Steel, M., Szekely, L. and Hendy, M. (1994). Reconstructing trees from sequences whose sites evolve at variable rates. *J. Comp. Biol.* **1**: 153–163.
- Steffansson, P. (2004). *Inferring Duplication and Loss Events Using Soft Parsimony*. MSc Thesis, Royal Institute of Technology, Stockholm, Sweden.
- Stevens, P.F. (1984). Homology and phylogeny: Morphology and systematics. *Syst. Bot.* **9**: 395–409.
- Stevens, P.F. (1991). Character states, Morphological variation, and phylogenetic analysis: A review. *Syst. Bot.* **16**: 553–583.
- Storm, C.E. and Sonnhammer, E.L. (2003). Comprehensive analysis of orthologous protein domains using the HOPS database. *Genome Res.* **13**: 2353–2362.
- Strong, E. and Lipscomb, D. (1999). Character coding and inapplicable data. *Cladistics* **15**: 363–371.
- Stuart, J.M., Segal, E., Koller, D. and Kim, S.K. (2003). A gene-coexpression network for global discovery of conserved genetic modules. *Science* **302**: 249–255.
- Suzuki, Y., Glazko, G. and Nei, M. (2002). Overcredibility of molecular phylogenetics obtained by Bayesian phylogenetics. *Proc. Natl. Acad. Sci. USA* **99**: 16138–16143.
- Swofford, D.L. (1984). *PAUP: Phylogenetic Analysis Using Parsimony*. Champaign, IL, Illinois Natural History Survey.
- Swofford, D.L. (1985). *PAUP: Phylogenetic Analysis Using Parsimony*, vers. 2.4. Champaign, IL, Illinois Natural History Survey.
- Swofford, D.L. (1990). *PAUP: Phylogenetic Analysis Using Parsimony*, vers. 3.0. (incl. vers. 3.0s). Champaign, Illinois Natural History Survey.
- Swofford, D.L. (1991). When are phylogeny estimates from molecular and morphological data incongruent? In *Phylogenetic Analysis of DNA Sequences*. (eds M.M. Miyamoto and J. Cracraft), pp. 295–333. New York, Oxford University Press.
- Swofford, D.L. (1993). *PAUP: Phylogenetic Analysis Using Parsimony*, vers. 3.1 (incl. vers. 3.1.1). Champaign, IL, Illinois Natural History Survey.

- Swofford, D.L. (2002). *PAUP*: Phylogenetic Analysis Using Parsimony (*and other methods)*, vers. 4 (incl. vers. 4.0b10). Sunderland, MA, Sinauer Associates.
- Swofford, D.L., Olsen, G.J., Waddell, P.J. and Hillis, D.M. (1996). Phylogenetic inference. In *Molecular Systematics*, 2nd edn (eds D.M. Hillis, C. Moritz and B.K. Marble), pp. 407–514. Sunderland, MA Sinauer Associates.
- Swofford, D., Waddell, P., Huelsenbeck, J., Foster, P., Lewis, P. and Rogers, J. (2001). Bias in phylogenetic estimation and its relevance to the choice between parsimony and likelihoods methods. *Syst. Biol.* **50**: 525–539.
- Tatusov, R.L., Koonin, E.V. and Lipman, D.J. (1997). A genomic perspective on protein families. *Science* **278**: 631–637.
- Tatusov, R.L., Fedorova, N.D., Jackson, J.D., Jacobs, A.R., Kiryutin, B., Koonin, E.V., Krylov, D.M., Mazumder, R., Mekhedov, S.L., Nikolskaya, A.N., *et al.* (2003). The COG database: An updated version includes eukaryotes. *BMC Bioinformatics* **4**: 41.
- Tehler, A., Little, D.P. and Farris, J.S. (2003). The full-length phylogenetic tree from 1 551 ribosomal sequences of chitinous fungi. *Fungi. Mycol. Res.* **107**: 901–916.
- Tellgren, Å., Berglund, A.C., Savolainen, P., Janis, C.M. and Liberles, D.A. (2004). Myostatin rapid sequence evolution in ruminants predates domestication. *Mol. Phylogenet. Evol.* **33**: 782–790.
- Thanaraj, T.A., Stamm, S., Clark, F., Riethoven, J.J., Le Texier, V. and Muilu, J. (2004). ASD: The Alternative Splicing Database. *Nucleic Acids Res.* **32**: D64–D69.
- Thayer, H.S. (1953). *Newton's Philosophy of Nature: Selections from his Writings*. New York, Hafner Publishing Co.
- Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994). CLUSTAL W: Improving the sensitivity of progressive multiple alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**: 4673–4680.
- Thorne, J.L., Kishino, H. and Felsenstein, J. (1991). An evolutionary model for maximum likelihood alignment of DNA sequences. *J. Mol. Evol.* **33**: 114–124.
- Thorne, J.L., Kishino, H. and Felsenstein, J. (1992). Inching toward reality: An improved likelihood model of sequence evolution. *J. Mol. Evol.* **34**: 3–16.
- Tierney, L. (1994). Markov chains for exploring posterior distributions. *Ann. Stat.* **22**: 1701–1786.
- Tuffley, C. and Steel, M. (1997). Links between maximum likelihood and maximum parsimony under a simple model of site substitution. *Bull. Math. Biol.* **59**: 581–607.
- Uddin, M., Wildman, D.E., Liu, G., Xu, W., Johnson, R.M., Hof, P.R., Kapatos, G., Grossman, L.I. and Goodman, M. (2004). Sister grouping of chimpanzees and humans as revealed by genome-wide phylogenetic analysis of brain gene expression profiles. *Proc. Natl. Acad. Sci. USA.* **101**: 2957–2962.
- Vander Stappen, J., De Laet, J., Gama-López, S., Van Campenhout, S. and Volckaert, G. (2002). Phylogenetic analysis of *Stylosanthes* (Fabaceae) based on the internal transcribed spacer region (ITS) of nuclear ribosomal DNA. *Plant Syst. Evol.* **234**: 27–51.
- Vingron, M. (1999). Sequence alignment and phylogeny construction. In *Mathematical Support for Molecular Biology* (eds M. Farach-Colton, F.S. Roberts, M. Vingron and M. Waterman), pp. 53–64. DIMACS Series in Discrete Mathematics and Theoretical Computer Science, vol. 47. Providence, RI, American Mathematical Society.
- Vrana, P. and Wheeler, W. (1992). Individual organisms as terminal entities: Laying the species problem to rest. *Cladistics* **8**: 67–72.
- Wagner, Jr., W.H. (1952). The fern genus *Diellia*: structure, affinities, and taxonomy. *Univ. Cal. Publ. Bot.* **26**: 1–212, pl. 1–21.
- Wagner, Jr., W.H. (1961). Problems in the classification of ferns. In *Recent Advances in Botany*, vol. 1, pp. 841–844. Toronto, University of Toronto Press.
- Walsh, D. (1979). Occam's razor: A principle of intellectual elegance. *Am. Phil. Q.* **16**: 241–244.
- Wang L. and Jiang, T. (1994). On the complexity of multiple sequence alignment. *J. Comput. Biol.* **1**: 337–348.
- Wang, L., Jiang, T. and Lawler, L. (1996). Approximation algorithms for tree alignment with a given phylogeny. *Algorithmica* **16**: 302–315.
- Wang, L.S., Jansen, R., Moret, B., Raubeson, L. and Warnow, T. (2002). Fast phylogenetic methods for the analysis of genome rearrangement data: An empirical study. *Proceedings of the Pacific Symposium on Biocomputing* (PSB 02), pp. 524–535. Singapore, World Scientific.
- Watanabe, H., Mori, H., Itoh, T. and Gojobori, T. (1997). Genome plasticity as a paradigm of eubacteria evolution. *J. Mol. Evol.* **44**: S57–S64.
- Waterston, R.H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J.F., Agarwal, P., Agarwala, R., Ainscough, R., Alexandersson, M., An, P. *et al.* (2002). Initial sequencing and comparative analysis of the mouse genome. *Nature* **420**: 520–562.
- Wheeler, Q.D. (1986). Character weighting and cladistic analysis. *Syst. Zool.* **35**: 102–109.
- Wheeler, Q.D. and Meier, R. (eds) (2000). *Species Concepts and Phylogenetic Theory: a Debate*, pp. 179–184. New York, Columbia University Press.
- Wheeler, W.C. (1994). Sources of ambiguity in nucleic acid sequence alignment. In *Molecular Ecology and Evolution: Approaches and Applications* (eds B. Schierwater,

- B. Streit, G.P. Wagner and R. DeSalle), pp. 323–352. Basel, Birkhäuser Verlag.
- Wheeler, W.C. (1996). Optimization alignment: The end of multiple sequence alignment in phylogenetics? *Cladistics* **12**: 1–9.
- Wheeler, W.C. (1998). Alignment characters, dynamic programming and heuristic solutions. In *Molecular Approaches to Ecology and Evolution* (eds R. DeSalle and B. Schierwater), pp. 243–251. Basel, Birkhäuser Verlag.
- Wheeler, W.C. (1999). Fixed character states and the optimization of molecular sequence data. *Cladistics* **15**: 379–385.
- Wheeler, W.C. (2001a). Homology and the optimization of DNA sequence data. *Cladistics* **17**: S3–S11.
- Wheeler, W. (2001b). Homology and DNA sequence data. In *The Character Concept in Evolutionary Biology* (ed. G.P. Wagner), pp. 303–317. San Diego, Academic Press.
- Wheeler, W.C. (2002). Optimization Alignment: Down, up, error, and improvements. In *Techniques in Molecular Systematics and Evolution* (eds R. DeSalle, G. Giribet and W. Wheeler), pp. 55–69. Basel, Birkhäuser Verlag.
- Wheeler, W.C. (2003a). Implied alignment: A synapomorphy-based multiple-sequence alignment method and its use in cladogram search. *Cladistics* **19**: 261–268.
- Wheeler, W.C. (2003b). Search-based optimization. *Cladistics*, **19**: 348–355.
- Wheeler, W.C. (2003c). Iterative pass optimization of sequence data. *Cladistics* **19**: 254–260.
- Wheeler, W.C. and Gladstein, D.S. (1994). MALIGN: A multiple sequence alignment program. *J. Hered.* **85**: 417–418.
- Wheeler, W.C. and Hayashi, C.Y. (1998). The phylogeny of the extant chelicerate orders. *Cladistics* **14**: 173–192.
- Wheeler, W., Gladstein, D. and De Laet, J. (2003). POY, ver. 3.0.11. Available at <ftp://ftp.amah.org/pub/molecular/poy>.
- Wiley, E.O. (1975). Karl R. Popper, systematics, and classification: A reply to Walter Bock and other evolutionary taxonomists. *Syst. Zool.* **24**: 233–243.
- Wiley, E.O. (1981). *Phylogenetics: the Theory and Practice of Phylogenetic Systematics*. New York, John Wiley and Sons.
- Wilkinson, M. (1995). A comparison of two methods of character construction. *Cladistics* **11**: 297–308.
- Wolf, Y.I., Rogozin, I.B., Grishin, N.V., Tatusov, R.L. and Koonin, E.V. (2001). Genome trees constructed using five different approaches suggest new major bacterial clades. *BMC Evol. Biol.* **1**: 8.
- Wolf, Y.I., Rogozin, I.B., Grishin, N.V. and Koonin, E.V. (2002). Genome trees and the tree of life. *Trends Genet.* **18**: 472–479.
- Wolf, Y.I., Rogozin, I.B. and Koonin, E.V. (2004). Coelomata and not Ecdysozoa: evidence from genome-wide phylogenetic analysis. *Genome Res.* **14**: 29–36.
- Wolfe, K.H. and Sharp, P.M. (1993). Mammalian gene evolution: Nucleotide sequence divergence between mouse and rat. *J. Mol. Evol.* **37**: 441–456.
- Woodger, J.H. (1929). *Biological principles: A critical study*. New York, Harcourt, Brace and Co.
- Wrinch, D. and Jeffreys, H. (1921). On certain fundamental principles of scientific inquiry. *Phil. Mag.* **42**: 369–390.
- Yang, Z. (1994). Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: Approximate methods. *J. Mol. Evol.* **39**: 306–314.
- Yang, Z. (1996). Phylogenetic analysis using parsimony and likelihood methods. *J. Mol. Evol.* **39**: 294–307.
- Yang, Z.H. (1998). Likelihood ratio tests for detecting positive selection and application to primate lysozyme evolution. *Mol. Biol. Evol.* **15**: 568–573.
- Yang, Z.H. and Bielawski, B. (2000). Statistical methods for detecting molecular adaptation. *Trends Ecol. Evol.* **15**: 496–503.
- Yang, Z. and Rannala, B. (1997). Bayesian phylogenetic inference using DNA sequences: A Markov chain Monte Carlo method. *Mol. Biol. Evol.* **14**: 717–724.
- Yang, Z., Goldman, N. and Friday, A. (1995a). Maximum likelihood trees from DNA sequences: A peculiar statistical estimation problem. *Syst. Biol.* **44**: 384–399.
- Yang, Z.H., Kumar, S. and Nei, M. (1995b). A new method of inference of ancestral nucleotide and amino acid sequences. *Genetics* **141**: 1641–1650.
- Yeates, D. (1992). Why remove autapomorphies? *Cladistics* **8**: 387–389.